

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



534 426

(43) International Publication Date  
17 June 2004 (17.06.2004)

PCT

(10) International Publication Number  
**WO 2004/050602 A1**

(51) International Patent Classification<sup>7</sup>: **C07C 69/28**,  
69/74, 69/66, A61K 7/46, C11B 9/00

(21) International Application Number:  
PCT/CH2003/000772

(22) International Filing Date:  
24 November 2003 (24.11.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0227807.5 29 November 2002 (29.11.2002) GB

(71) Applicant (for all designated States except US): **GIVAUDAN SA** [CH/CH]; Chemin de la Parfumerie 5, CH-1214 Vernier (CH).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **KRAFT, Philip** [DE/CH]; Kirchbachstrasse 4, CH-8600 Dübendorf (CH).

(74) Agent: **McSTEa, John, Anthony**; Givaudan Schweiz AG, Global Patents, Überlandstrasse 138, CH-8600 Dübendorf (CH).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ESTERS AND THEIR USE IN PERFUMERY



(I)

(57) Abstract: This invention relates to unsaturated alicyclic carbonyl compounds of formula (I) [insert formula (I) here] wherein R is C1 to C4 alkyl; or R is vinyl or a linear, branched or cyclic C3 to C4 alkenyl; X is carbonyl or a divalent radical -(CMe2)-; and Y is oxygen or a divalent radical -(CH2)-.

WO 2004/050602 A1

## ESTERS AND THEIR USE IN PERFUMERY

This invention relates to new odourant compounds having musk characteristics, their manufacture and their use in fragrance compositions.


5

Conventional compounds having musk characteristics have been selected from nitro arenes, polycyclic aromatics and macrocyclic compounds. However, in recent years there has been great activity to find novel compounds having musk characteristics to replace these conventional musks, the use of which is becoming more restricted because of, e.g. environmental concerns.


10

In recent years, research activity has resulted in the development of new classes of compounds with musk characteristics. EP 472966, for example describes a family of compounds exemplified by the product Helvetolide (1) that is described as having musky, ambrette-like characteristics. Further attempts were made to improve on the olfactory properties of Helvetolide (1) and its related compounds by replacing the gem-dialkyl group in the aliphatic side chain by a carbonyl group, as disclosed in WO 00/14051, exemplified by compound (2) that has been described as having a stronger musky smell than prior art compounds.

20



Helvetolide (1)




Compound (2)

Surprisingly, we now found certain unsaturated alicyclic carbonyl compounds that have musk characteristics and a high impact in perfume formulations.

Thus, the present invention refers in a first aspect to a compound of formula (I)

30

2



(I)

wherein

R is C<sub>1</sub> to C<sub>4</sub> alkyl, for example methyl, ethyl, *i*-propyl, *n*-propyl, *n*-butyl, *sec*-butyl, *tert*-butyl, cyclopropyl, cyclobutyl; or

R is vinyl or a linear, branched or cyclic C<sub>3</sub> to C<sub>4</sub> alkenyl, for example propen-1-yl, propen-2-yl, or prop-2-en-1-yl, butenyl, e.g. but-1-en-1-yl, cyclobut-1-en-1-yl, or butadienyl, e.g. buta-2,4-dien-1-yl;

X is carbonyl or a divalent radical  $-(CMe_2)-$  ;

Y is oxygen or a divalent radical  $-(CH_2)-$  ;

the bond between C-a and C-b is a single bond and the bond between C-b and C-c together with the dotted line represents a double bond; or

the bond between C-a and C-b together with the dotted line represents a double bond and the bond between C-b and C-c is a single bond.

The compounds according to the present invention may comprise one or more chiral centres and as such may exist as a mixture of stereoisomers, or they may be resolved as isomerically pure forms. Resolving stereoisomers adds to the complexity of manufacture and purification of these compounds and so it is preferred to use the compounds as mixtures of their stereoisomers simply for economic reasons. However, if it is desired to prepare individual stereoisomers, this may be achieved according to methodology known in the art, e.g. preparative HPLC and GC or by stereoselective syntheses.

Preferred compounds of formula (I) are propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, cyclopropanecarboxylic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, propionic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, cyclopropanecarboxylic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, propionic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester, and

cyclopropanecarboxylic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy-carbonylmethyl ester.

Particular preferred are compounds of formula (I) wherein the bond between C-b and C-  
5 c together with the dotted line represents a double bond and the bond between C-a and C-b is a single bond, e.g. propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester and cyclopropanecarboxylic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester.

10 Compounds of formula (I) having a double bond between C-b and C-c are more powerful and possess more distinct musk notes, while compounds having a double bond between C-a and C-b possess more pronounced fruity, green aspects besides their main musk character.

15 The compounds according to the present invention may be used alone or in combination with known odourant molecules selected from the extensive range of natural and synthetic molecules currently available, such as essential oils, alcohols, aldehydes and ketones, ethers and acetals, esters and lactones, macrocycles and heterocycles, and/or in admixture with one or more ingredients or excipients  
20 conventionally used in conjunction with odourants in fragrance compositions, for example carrier materials, and other auxiliary agents commonly used in the art.

The following list comprises examples of known odourant molecules, which may be combined with the compounds of the present invention:

25

– ethereal oils and extracts, e.g. castoreum, costus root oil, oak moss absolute, geranium oil, jasmin absolute, patchouli oil, rose oil, sandalwood oil or ylang-ylang oil;

30

– alcohols, e.g. citronellol, Ebanol<sup>TM</sup>, eugenol, geraniol, Super Muguet<sup>TM</sup>, linalool, phenylethyl alcohol, Sandalore<sup>TM</sup>, terpineol or Timberol<sup>TM</sup>.

– aldehydes and ketones, e.g.  $\alpha$ -amylcinnamaldehyd, Georgywood<sup>TM</sup>, hydroxycitronellal, Iso E Super<sup>TM</sup>, Isoraldeine<sup>TM</sup>, Hedione<sup>TM</sup>, maltol, methyl cedryl ketone, methylionone or vanillin;

35

- ether and acetals, e.g. Ambrox™, geranyl methyl ether, rose oxide or Spirambrene™.
- esters and lactones, e.g. benzyl acetat, cedryl acetate,  $\gamma$ -decalactone, Helvetolide® (1),  $\gamma$ -undecalactone or vetiveryl acetate.
- macrocycles, e.g. ambrettolide, ethylene brassylate or Exaltolide®.
- heterocycles, e.g. isobutylcholine.

10 However, due to their unique character, the compounds of formula (I) are especially well suited for use in fresh musky accords, woody-spicy or floral-hesperidic compositions as is more specifically illustrated in the Example.

The compounds of the present invention may be used in a broad range of fragrance applications, e.g. in any field of fine and functional perfumery, such as perfumes, household products, laundry products, body care products and cosmetics. The compounds can be employed in wide ranging amounts depending upon the specific application and on the nature and quantity of other odourant ingredients, that may be for example, from about 0.001 to about 20 weight percent. In one embodiment compounds of the present invention may be employed in a fabric softener in an amount of about 0.001 to 0.05 weight percent. In another embodiment compounds of the present invention may be used in an alcoholic solution in amounts of about 0.1 to 20 weight percent, more preferably between about 0.1 and 5 weight percent. However, these values should not be limiting on the present invention, since the experienced perfumer may also achieve effects or may create novel accords with lower or higher concentrations.

The compounds of the present invention may be employed into the fragrance application simply by direct mixing the fragrance composition with the fragrance application, or they may, in an earlier step be entrapped with an entrapment material such as for example polymers, capsules, microcapsules and nanocapsules, liposomes, film formers, absorbents such as for example by using carbon or zeolites, cyclic oligosaccharides and mixtures thereof, or they may be chemically bound to substrates which are adapted to release the fragrance molecule upon application of an exogenous stimulus such as light, enzyme, or the like, and then mixed with the application.

Compounds of formula (I) may be prepared starting from a corresponding allylic alcohol, which is accessible either by reduction of Artemone (1-(3',3'-dimethylcyclohex-1'-enyl)ethanone; trademark of Givaudan SA, Switzerland), or by Rupe rearrangement of 1-ethynyl-3,3-dimethylcyclohexanol.

By etherification of the corresponding allylic alcohol with isobutylene oxide and subsequent esterification with the corresponding carboxylic acids compounds of formula (I) wherein X is a divalent radical  $-(CMe_2)-$ , and Y is oxygen, i.e. oxa esters, may be synthesized.

Compounds of formula (I) wherein X is carbonyl and Y oxygen, i.e. diester, may be synthesized by esterification of the corresponding allylic alcohol with chloroacetic acid, followed by further esterification with the corresponding carboxylic acids.

Compounds of formula (I) wherein X is carbonyl and Y is a divalent radical  $-(CH_2)-$ , i.e. oxo ester, may be prepared by esterification of the corresponding allylic alcohol with the corresponding oxo carboxylic acids, e.g. laevulinic acid.

Compounds of formula (I) wherein X is a divalent radical  $-(CMe_2)-$  and Y is a divalent radical  $-(CH_2)-$ , i.e. oxa ketones may be prepared by etherification of the corresponding allylic alcohol with isobutylene oxide, subsequent oxidation to the aldehyde followed by a Wittig-Horner-Emmons reaction well known in the art and selective hydrogenation of the formed double bond.

Further particulars as to reaction conditions are provided in the examples.

There now follows a series of examples that illustrate the invention.

Example 1: Propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methyl-propyl ester

A solution of Artemone [1-(3',3'-dimethylcyclohex-1'-enyl)ethanone, commercial product of Givaudan, 152 g, 1.00 mol] in Et<sub>2</sub>O (500 ml) was added dropwise with stirring within 3 h at room temp. to a suspension of lithium aluminum hydride (LAH, 10.4 g, 275 mmol)

in Et<sub>2</sub>O (1 L). The reaction mixture was heated to reflux for 150 min., and after cooling down to 0° C quenched by careful dropwise addition of water (50 ml). Then 2N aq. HCl (200 ml) was added, and the mixture was poured into water (200 ml). The product was extracted with Et<sub>2</sub>O (2× 500 ml), and the combined extracts were washed with water (200 ml) and brine (100 ml), dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to dryness. The resulting residue (154 g) was purified by silica-gel FC (pentane/Et<sub>2</sub>O, 4:1) to afford 133 g (87%) of 1-(3',3'-dimethylcyclohex-1'-enyl)ethanol.

During a period of 1h, a 1M solution of MeAlCl<sub>2</sub> (150 ml, 150 mmol) in hexane was added dropwise with stirring at 0 °C to a solution of 1-(3',3'-dimethylcyclohex-1'-enyl)ethanol (46.3 g, 300 mmol) and isobutylene oxide (26.0 g, 360 mmol) in cyclohexane (300 ml). The cooling bath was removed, and stirring was continued at room temp. for 20 h, prior to pouring the reaction mixture into ice/water (1:1, 200 ml). Conc. aq. H<sub>3</sub>PO<sub>4</sub> was added until the slurry dissolved, and the product was extracted with Et<sub>2</sub>O (2× 200 ml). The combined organic extracts were washed with water (200 ml) and brine (25 ml), dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated in a rotary evaporator. The resulting residue (60.5 g) was purified by silica-gel FC (pentane/Et<sub>2</sub>O, 9:1, *R<sub>f</sub>* = 0.14) followed by distillation at 55°C /1.5 mbar to furnish 12.7 (19%) of 2-[1'-(3'',3''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol.

At 0°C under N<sub>2</sub>, *N,N'*-Dicyclohexylcarbodiimide (DCC, 2.27 g, 11.0 mmol) was added to a stirred solution of 2-[1'-(3'',3''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol (2.26 g, 10.0 mmol), propionic acid (740 mg, 10.0 mmol) and 4-(dimethylamino)pyridine (DMAP, 120 mg, 1.00 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (15 ml). The cooling bath was removed and the reaction mixture was stirred for 2 h at room temp. prior to vacuum filtration of the precipitate. The precipitate was washed with CH<sub>2</sub>Cl<sub>2</sub> (2×), and the combined filtrates were concentrated under reduced pressure. The crude material (3.25 g) was purified by silica-gel FC (pentane/Et<sub>2</sub>O, 19:1, *R<sub>f</sub>* = 0.46) to afford 2.52 g (89%) of the odoriferous title compound.

IR (ATR):  $\nu$  = 1741 cm<sup>-1</sup> (s,  $\nu$ O=CO), 1168 / 1068 cm<sup>-1</sup> (s,  $\nu$ C—O), 1366 cm<sup>-1</sup> (m,  $\delta$ CH<sub>3</sub>).— <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 0.93 / 0.94 (2s, 6H, 3'''-Me<sub>2</sub>), 1.14 (d, *J* = 6.5 Hz, 3H, 2''-H<sub>3</sub>), 1.16 (t, *J* = 7.5 Hz, 3H, 3-H<sub>3</sub>), 1.17 / 1.18 (2s, 6H, 2'-Me<sub>2</sub>), 1.37 (m<sub>c</sub>, 2H, 4'''-H<sub>2</sub>), 1.60 (m<sub>c</sub>, 2H, 5'''-H<sub>2</sub>), 1.81–2.06 (m, 2H, 6'''-H<sub>2</sub>), 2.37 (q, *J* = 7.5 Hz, 2H, 2-H<sub>2</sub>), 3.90 (d,

$J = 11.0$  Hz, 1H, 1'-H<sub>b</sub>), 3.99 (q,  $J = 7.5$  Hz, 1H, 1''-H), 4.01 (d,  $J = 11.0$  Hz, 1H, 1'-H<sub>a</sub>), 5.30 (s, 1H, 2'''-H). –  $^{13}\text{C}$  NMR (CDCl<sub>3</sub>):  $\delta = 8.99$  (q, C-3), 19.6 (t, C-5'''), 22.4 (q, C-2''), 23.4 (t, C-6'''), 23.5 / 23.6 (2q, 2'-Me<sub>2</sub>), 27.5 (t, C-2), 29.3 / 29.9 (2q, 3'''-Me<sub>2</sub>), 31.2 (s, C-3'''), 37.3 (t, C-4'''), 69.7 (t, C-1'), 72.2 (d, C-1''), 74.3 (s, C-2'), 131.3 (d, C-2'''), 139.0 (s, C-1'''), 174.1 (s, C-1). – MS (70 eV);  $m/z = 153$  (15) [C<sub>10</sub>H<sub>17</sub>O<sup>+</sup>], 147 (3) [C<sub>7</sub>H<sub>15</sub>O<sub>3</sub><sup>+</sup>], 137 (67) [C<sub>10</sub>H<sub>17</sub><sup>+</sup>], 129 (36) [C<sub>7</sub>H<sub>13</sub>O<sub>2</sub><sup>+</sup>], 121 (29) [C<sub>9</sub>H<sub>13</sub><sup>+</sup>], 107 (17) [C<sub>8</sub>H<sub>11</sub><sup>+</sup>], 95 (28) [C<sub>7</sub>H<sub>11</sub><sup>+</sup>], 93 (27) [C<sub>7</sub>H<sub>9</sub><sup>+</sup>], 79 (19) [C<sub>6</sub>H<sub>7</sub><sup>+</sup>], 57 (100) [C<sub>3</sub>H<sub>5</sub><sup>+</sup>].

Odor description: Musky, powerful, powdery, slightly animalic.

Example 2: Cyclopropanecarboxylic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)-ethoxy]-2'-methylpropyl ester

Following the same procedure according to Example 1, Steglich esterification of 2-[1'-(3'',3''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol (2.26 g, 10.0 mmol) with cyclopropanecarboxylic acid (860 mg, 10.0 mmol), and purification by silica-gel FC (pentane/Et<sub>2</sub>O, 19:1,  $R_f = 0.33$ ) furnished 2.68 (91%) of cyclopropanecarboxylic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester.

IR (ATR):  $\nu = 1731$  cm<sup>-1</sup> (s,  $\nu\text{O}=\text{CO}$ ), 1153 / 1068 cm<sup>-1</sup> (s,  $\nu\text{C}-\text{O}$ ), 1400 / 1381 / 1366 cm<sup>-1</sup> (m,  $\delta\text{CH}_3$ ). –  $^1\text{H}$  NMR (CDCl<sub>3</sub>):  $\delta = 0.87$  (m<sub>c</sub>, 2H, 3-,4-H<sub>b</sub>), 0.93 / 0.94 (2s, 6H, 3'''-Me<sub>2</sub>), 1.02 (m<sub>c</sub>, 2H, 3-,4-H<sub>a</sub>), 1.15 (d,  $J = 6.5$  Hz, 3H, 2''-H<sub>3</sub>), 1.17 / 1.18 (2s, 6H, 2'-Me<sub>2</sub>), 1.38 (m<sub>c</sub>, 2H, 4'''-H<sub>2</sub>), 1.62 (m<sub>c</sub>, 2H, 5'''-H<sub>2</sub>), 1.81–2.06 (m, 2H, 6'''-H<sub>2</sub>), 3.89 (d,  $J = 11.0$  Hz, 1H, 1'-H<sub>b</sub>), 3.98 (q,  $J = 7.5$  Hz, 1H, 1''-H), 4.00 (d,  $J = 11.0$  Hz, 1H, 1'-H<sub>a</sub>), 5.30 (s, 1H, 2'''-H). –  $^{13}\text{C}$  NMR (CDCl<sub>3</sub>):  $\delta = 8.18$  / 8.19 (2t, C-3,-4), 12.8 (d, C-2), 19.6 (t, C-5'''), 22.4 (q, C-2''), 23.4 (t, C-6'''), 23.5 (2q, 2'-Me<sub>2</sub>), 29.3 / 29.9 (2q, 3'''-Me<sub>2</sub>), 31.2 (s, C-3'''), 37.3 (t, C-4'''), 69.8 (t, C-1'), 72.2 (d, C-1''), 74.3 (s, C-2'), 131.3 (d, C-2'''), 139.0 (s, C-1'''), 174.5 (s, C-1). – MS (70 eV);  $m/z = 294$  (1) [M<sup>+</sup>], 279 (1) [M<sup>+</sup> – CH<sub>3</sub>], 159 (3) [C<sub>8</sub>H<sub>15</sub>O<sub>3</sub><sup>+</sup>], 153 (22) [C<sub>10</sub>H<sub>17</sub>O<sup>+</sup>], 141 (30) [C<sub>8</sub>H<sub>13</sub>O<sub>2</sub><sup>+</sup>], 137 (61) [C<sub>10</sub>H<sub>17</sub><sup>+</sup>], 121 (27) [C<sub>9</sub>H<sub>13</sub><sup>+</sup>], 107 (17) [C<sub>8</sub>H<sub>11</sub><sup>+</sup>], 95 (26) [C<sub>7</sub>H<sub>11</sub><sup>+</sup>], 93 (25) [C<sub>7</sub>H<sub>9</sub><sup>+</sup>], 81 (27) [C<sub>6</sub>H<sub>9</sub><sup>+</sup>], 79 (17) [C<sub>6</sub>H<sub>7</sub><sup>+</sup>], 69 (100) [C<sub>5</sub>H<sub>9</sub><sup>+</sup>].

Odor description: Musky, pleasant, powdery, strong.



Example 3: Propionic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester

Phosphorus pentoxide (33.1 g, 233 mmol) was added to a solution of 1-ethynyl-3,3-dimethylcyclohexanol (152 g, 1.00 mol) in MePh (800 ml). The slurry was heated to reflux, and stirred at this temp. for 90 min. The reaction mixture was allowed to cool to room temp., and then poured into ice/water (1:1, 500 ml). The product was extracted with Et<sub>2</sub>O (2× 500 ml), and the combined organic extracts were washed with water (500 ml) and brine (100 ml), dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure.

10 Silica-gel FC (pentane/ Et<sub>2</sub>O, 9:1, *R<sub>f</sub>* = 0.70) provided 13.8 g (9 %) of 1-(5',5'-dimethylcyclohex-1'-enyl)ethanone.

A solution of 1-(5',5'-dimethylcyclohex-1'-enyl)ethanone (13.1 g, 85.8 mmol) in Et<sub>2</sub>O (50 ml) was added dropwise with stirring within 50 min. to a suspension of LAH (895 mg, 23.6 mmol) in Et<sub>2</sub>O (150 ml). The reaction mixture was refluxed for 1 h prior to quenching at 0 °C by careful addition of water (50 ml) followed by 5 N aq. HCl (50 ml). The organic layer was separated and the aqueous one extracted with Et<sub>2</sub>O. The combined ethereal solutions were washed with water and brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated in a rotary evaporator. Silica-gel FC (pentane/Et<sub>2</sub>O, 9:1, *R<sub>f</sub>* = 0.14) of the

15 20 resulting residue (14.7 g) gave 11.2 g (85 %) of 1-(5',5'-dimethylcyclohex-1'-enyl)ethanol.

At 0 °C under N<sub>2</sub>, a 1 M solution of MeAlCl<sub>2</sub> (33.6 ml, 33.6 mmol) in hexane was added dropwise during 1 h to a stirred solution of 1-(5',5'-dimethyl-cyclohex-1'-enyl)ethanol (10.4 g, 67.2 mmol) and isobutylene oxide (5.82 g, 80.7 mmol) in cyclohexane (67 ml). The cooling bath was removed, the reaction mixture stirred at room temp. for 23 h, and then poured into ice/water (1:1, 200 ml). The slurry was brought into solution by addition of conc. aq. H<sub>3</sub>PO<sub>4</sub>, and the product was extracted with Et<sub>2</sub>O (2× 100 ml). The combined organic extracts were washed with water (100 ml) and brine (25 ml), dried

25 30 (Na<sub>2</sub>SO<sub>4</sub>), and concentrated in a rotary evaporator. The crude material (12.4 g) was purified by silica-gel FC (pentane/Et<sub>2</sub>O, 9:1, *R<sub>f</sub>* = 0.17) to provide 3.31 g (22 %) of 2-[1'-(5'',5''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol.

Following the same procedure according to Example 1, Steglich esterification of 2-[1'-(5'',5''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol (1.29 g, 5.70 mmol) with

35

propionic acid (420 mg, 5.70 mmol), and purification by silica-gel FC (pentane/Et<sub>2</sub>O, 9:1, *R<sub>f</sub>* = 0.56) furnished 420 mg (26%) of propionic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester.

- 5 IR (ATR):  $\nu$  = 1169 / 1068 cm<sup>-1</sup> (s,  $\nu$ C–O), 1741 cm<sup>-1</sup> (s,  $\nu$ O=CO), 1365 cm<sup>-1</sup> (m,  $\delta$ CH<sub>3</sub>).  
 – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 0.89 / 0.91 (2s, 6H, 5'''-Me<sub>2</sub>), 1.14 (d, *J* = 6.5 Hz, 3H, 2''-H<sub>3</sub>),  
 1.16 (t, *J* = 7.5 Hz, 3H, 3-H<sub>3</sub>), 1.17 / 1.18 (2s, 6H, 2'-Me<sub>2</sub>), 1.29 (t, *J* = 6.5 Hz, 2H, 4'''-  
 H<sub>2</sub>), 1.69 (dd, *J* = 17.0, 2.0 Hz, 1H, 6'''-H<sub>b</sub>), 1.83 (dd, *J* = 17.0, 2.0 Hz, 1H, 6'''-H<sub>a</sub>), 2.01  
 (m<sub>c</sub>, 2H, 3'''-H<sub>2</sub>), 2.36 (q, *J* = 7.5 Hz, 2H, 2-H<sub>2</sub>), 3.92 (d, *J* = 11.0 Hz, 1H, 1'-H<sub>b</sub>), 4.00 (d,  
 10 *J* = 11.0 Hz, 1H, 1'-H<sub>a</sub>), 4.02 (q, *J* = 6.5 Hz, 1''-H), 5.54 (s, 1H, 2'''-H). – <sup>13</sup>C NMR  
 (CDCl<sub>3</sub>):  $\delta$  = 8.98 (q, C-3), 22.4 (q, C-2''), 22.7 (t, C-3'''), 23.4 (t, C-6'''), 23.4 / 23.6 (2q,  
 2'-Me<sub>2</sub>), 27.5 (t, C-2), 27.9 / 28.0 (2q, 5'''-Me<sub>2</sub>), 28.6 (s, C-5'''), 35.2 (t, C-4'''), 37.5 (t, C-  
 6'''), 69.8 (t, C-1'), 71.9 (d, C-1''), 74.2 (s, C-2'), 119.1 (d, C-2'''), 140.4 (s, C-1'''), 174.1  
 (s, C-1). – MS (70 eV); *m/z* = 153 (16) [C<sub>10</sub>H<sub>17</sub>O<sup>+</sup>], 147 (2) [C<sub>7</sub>H<sub>15</sub>O<sub>3</sub><sup>+</sup>], 137 (59) [C<sub>10</sub>H<sub>17</sub><sup>+</sup>],  
 15 129 (30) [C<sub>7</sub>H<sub>13</sub>O<sub>2</sub><sup>+</sup>], 121 (37) [C<sub>9</sub>H<sub>13</sub><sup>+</sup>], 107 (29) [C<sub>8</sub>H<sub>11</sub><sup>+</sup>], 95 (29) [C<sub>7</sub>H<sub>11</sub><sup>+</sup>], 93 (39)  
 [C<sub>7</sub>H<sub>9</sub><sup>+</sup>], 79 (48) [C<sub>6</sub>H<sub>7</sub><sup>+</sup>], 57 (100) [C<sub>4</sub>H<sub>9</sub><sup>+</sup>].

Odor description: Musky, powdery, fruity.

20 Example 4: Cyclopropanecarboxylic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)-ethoxy]-2'-methylpropyl ester

- Following the same procedure according to Example 1, Steglich esterification of 2-[1'-  
 (5'',5''-dimethylcyclohex-1''-enyl)ethoxy]-2-methylpropan-1-ol (1.29 g, 5.70 mmol) with  
 25 cyclopropanecarboxylic acid (490 mg, 5.70 mmol), and purification by silica-gel FC  
 (pentane/Et<sub>2</sub>O, 9:1, *R<sub>f</sub>* = 0.58) furnished 590 mg (35%) of cyclopropanecarboxylic acid  
 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester.

- IR (ATR):  $\nu$  = 1163 / 1067 cm<sup>-1</sup> (s,  $\nu$ C–O), 1730 cm<sup>-1</sup> (s,  $\nu$ O=CO), 1382 / 1400 / 1382  
 30 cm<sup>-1</sup> (m,  $\delta$ CH<sub>3</sub>). – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 0.85 (dt, *J* = 8.0, 4.5 Hz, 2H, 3-,4-H<sub>b</sub>), 0.89 /  
 0.91 (s, 6H, 5'''-Me<sub>2</sub>), 1.01 (dt, *J* = 8.0, 4.5, 2H, 3-,4-H<sub>a</sub>), 1.14 (d, *J* = 6.5 Hz, 3H, 2''-H<sub>3</sub>),  
 1.17 / 1.18 (2s, 6H, 2'-Me<sub>2</sub>), 1.29 (dd, *J* = 6.5, 2.5 Hz, 1H, 4-H<sub>b</sub>), 1.30 (dd, *J* = 6.5, 2.5  
 Hz, 1H, 4-H<sub>a</sub>), 1.64 (tt, *J* = 8.0, 4.5 Hz, 1H, 2-H), 1.70 (dd, *J* = 17.0, 2.0 Hz, 1H, 6'''-H<sub>b</sub>),  
 1.84 (dd, *J* = 17.0, 2.0 Hz, 1H, 6'''-H<sub>a</sub>), 2.01 (m<sub>c</sub>, 2H, 3'''-H<sub>2</sub>), 3.91 (d, *J* = 11.0 Hz, 1H,

1'-H<sub>b</sub>), 3.98 (d, *J* = 11.0 Hz, 1H 1'-H<sub>a</sub>), 4.02 (q, *J* = 6.5 Hz, 1H, 1''-H), 5.54 (s, 1H, 2'''-H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 8.14 (2t, C-3,-4), 12.8 (d, C-2), 22.4 (q, C-2''), 22.7 (t, C-3'''), 23.5 / 23.6 (2q, 2'-Me<sub>2</sub>), 27.9 / 28.0 (2q, 5'''-Me<sub>2</sub>), 28.6 (s, C-5'''), 35.2 (t, C-4'''), 37.5 (t, C-6'''), 69.8 (t, C-1'), 71.9 (d, C-1''), 74.3 (s, C-2'), 119.1 (d, C-2'''), 140.3 (s, C-1'''), 174.5 (s, C-1). – MS (70 eV); *m/z* = 294 (1) [M<sup>+</sup>], 159 (2) [C<sub>8</sub>H<sub>15</sub>O<sub>3</sub><sup>+</sup>], 153 (25) [C<sub>10</sub>H<sub>17</sub>O<sup>+</sup>], 141 (27) [C<sub>8</sub>H<sub>13</sub>O<sub>2</sub><sup>+</sup>], 137 (54) [C<sub>10</sub>H<sub>17</sub><sup>+</sup>], 121 (27) [C<sub>9</sub>H<sub>13</sub><sup>+</sup>], 107 (22) [C<sub>8</sub>H<sub>11</sub><sup>+</sup>], 95 (25) [C<sub>7</sub>H<sub>11</sub><sup>+</sup>], 93 (26) [C<sub>7</sub>H<sub>9</sub><sup>+</sup>], 81 (19) [C<sub>6</sub>H<sub>9</sub><sup>+</sup>], 79 (30) [C<sub>6</sub>H<sub>7</sub><sup>+</sup>], 69 (100) [C<sub>5</sub>H<sub>9</sub><sup>+</sup>].

Odor description: Musky, fresh, floral, slightly metallic.

Example 5: Propionic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester

*N,N'*-Dicyclohexylcarbodiimide (DCC, 2.58 g, 12.5 mmol) was added at 0 °C under N<sub>2</sub> to a solution of 1-(5,5-dimethylcyclohex-1-enyl)ethanol, chloroacetic acid (1.07 g, 11.3 mmol) and 4-(dimethylamino)pyridine (DMAP, 140 mg, 1.13 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (15 ml). The cooling bath was removed and the reaction mixture was stirred for 1 h at room temp., prior to vacuum filtration of the precipitates. The filtrate was concentrated under reduced pressure, the resulting residue purified by silica-gel FC (pentane/Et<sub>2</sub>O, 19:1, *R<sub>f</sub>* = 0.65) to furnish 2.17 g (83 %) of chloroacetic acid 1'-(5'',5''-dimethylcyclohex-1''-enyl)ethyl ester.

A mixture of chloroacetic acid 1'-(5'',5''-dimethylcyclohex-1''-enyl)ethyl ester (1.00 g, 4.33 mmol), propionic acid (320 mg, 4.33 mmol), K<sub>2</sub>CO<sub>3</sub> (1.20 g, 8.67 mmol) and NaBr (450 mg, 4.33 mmol) in Et<sub>2</sub>CO / dioxane (4:1, 10 ml) was refluxed for 1 day prior to pouring into water (50 ml). The product was extracted with Et<sub>2</sub>O (2× 50 ml), and the combined extracts were washed with water (50 ml) and brine (25 ml). After drying with Na<sub>2</sub>SO<sub>4</sub> and evaporation of the solvent under reduced pressure, silica-gel FC (pentane/Et<sub>2</sub>O, 9:1, *R<sub>f</sub>* = 0.41) afforded 370 mg (32%) of propionic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester.

IR (ATR): ν = 1161 cm<sup>-1</sup> (s, νC—O), 1747 cm<sup>-1</sup> (s, νO=CO). – <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 0.89 / 0.91 (2s, 6H, 5'''-Me<sub>2</sub>), 1.19 (t, *J* = 7.5 Hz, 3H, 3-H, 3-H<sub>3</sub>), 1.31 (d, *J* = 6.5 Hz, 3H, 2''-H<sub>3</sub>), 1.34 (m<sub>c</sub>, 2H, 4'''-H<sub>2</sub>), 1.69 (d, *J* = 16.5 Hz, 1H, 6'''-H<sub>b</sub>), 1.78 (d, *J* = 16.5 Hz, 1H, 6'''-H<sub>a</sub>), 2.05 (m<sub>c</sub>, 2H, 3'''-H<sub>2</sub>), 2.45 (q, *J* = 7.5 Hz, 2H, 2-H<sub>2</sub>), 4.56 (d, *J* = 16.0 Hz, 1H,

2'-H<sub>b</sub>), 4.61 (d,  $J = 16.0$  Hz, 1H, 2'-H<sub>a</sub>), 5.32 (q,  $J = 6.5$  Hz, 1H, 1''-H), 5.67 (s, 1H, 2'''-H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 8.81$  (q, C-3), 18.5 (q, C-2''), 22.7 (t, C-3'''), 27.0 (t, C-2), 27.5 / 28.3 (2q, 5'''-Me<sub>2</sub>), 28.6 (s, C-5'''), 34.7 (t, C-4'''), 37.5 (t, C-6'''), 60.6 (t, C-2'), 75.5 (d, C-1''), 123.1 (d, C-2'''), 135.2 (s, C-1'''), 167.1 (s, C-1'), 173.5 (s, C-1). – MS (70 eV);  
 5  $m/z = 154$  (3) [C<sub>10</sub>H<sub>18</sub>O<sup>+</sup>], 136 (58) [C<sub>10</sub>H<sub>16</sub><sup>+</sup>], 121 (86) [C<sub>9</sub>H<sub>13</sub><sup>+</sup>], 107 (75) [C<sub>8</sub>H<sub>11</sub><sup>+</sup>], 93 (100) [C<sub>7</sub>H<sub>9</sub><sup>+</sup>], 79 (100) [C<sub>6</sub>H<sub>7</sub><sup>+</sup>], 41 (36) [C<sub>3</sub>H<sub>5</sub><sup>+</sup>].

Odor description: Musky, green.

10 Example 6: Cyclopropanecarboxylic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy-carbonylmethyl ester

A mixture of chloroacetic acid 1'-(5'',5''-dimethylcyclohex-1''-enyl)ethyl ester (1.00 g, 4.33 mmol), cyclopropanecarboxylic acid (370 mg, 4.33 mmol), K<sub>2</sub>CO<sub>3</sub> (1.20 g, 8.67  
 15 mmol) and NaBr (450 mg, 4.33 mmol) in Et<sub>2</sub>CO / dioxane (4:1, 10 ml) was refluxed for 1 day prior to pouring into water (50 ml). The product was extracted with Et<sub>2</sub>O (2× 50 ml), and the combined extracts were washed with water (50 ml) and brine (25 ml). After drying with Na<sub>2</sub>SO<sub>4</sub> and evaporation of the solvent under reduced pressure, silica-gel FC (pentane/Et<sub>2</sub>O, 9:1,  $R_f = 0.30$ ) afforded 870 mg (72%) of cyclopropanecarboxylic  
 20 acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester.

IR (ATR):  $\nu = 1156$  cm<sup>-1</sup> (s,  $\nu$ C–O), 1736 cm<sup>-1</sup> (s,  $\nu$ O=CO). – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 0.89 / 0.91$  (2s, 6H, 5'''-Me<sub>2</sub>), 0.93 (m<sub>c</sub>, 2H, 3-,4-H<sub>b</sub>), 1.07 (m<sub>c</sub>, 2H, 3-,4-H<sub>a</sub>), 1.30 (d,  $J = 6.5$  Hz, 3H, 2''-H<sub>3</sub>), 1.33 (m<sub>c</sub>, 2H, 4'''-H<sub>2</sub>), 1.68–1.80 (m, 3H, 2-H, 6'''-H<sub>2</sub>), 2.05 (m<sub>c</sub>, 2H, 3'''-H<sub>2</sub>), 4.56 (d,  $J = 16.0$  Hz, 1H, 2'-H<sub>b</sub>), 4.61 (d,  $J = 16.0$  Hz, 1H, 2'-H<sub>a</sub>), 5.30 (q,  $J = 6.5$  Hz, 1H, 1''-H), 5.67 (s, 1H, 2'''-H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 8.67 / 8.65$  (t, C-3,-4), 12.4 (d, C.-2), 18.5 (q, C-2''), 22.7 (t, C-3'''), 27.5 / 28.3 (2q, 5'''-Me<sub>2</sub>), 28.5 (s, C-5'''), 34.7 (t, C-4'''), 37.5 (t, C-6'''), 60.6 (t, C-2'), 75.5 (d, C-1''), 123.0 (d, C-2'''), 135.2 (s, C-1'''), 167.1 (s, C-1'), 174.0 (s, C-1). – MS (70 eV);  $m/z = 153$  (2) [C<sub>10</sub>H<sub>17</sub>O<sup>+</sup>], 136 (61) [C<sub>10</sub>H<sub>16</sub><sup>+</sup>], 121 (84) [C<sub>9</sub>H<sub>13</sub><sup>+</sup>], 107 (71) [C<sub>8</sub>H<sub>11</sub><sup>+</sup>], 93 (98) [C<sub>7</sub>H<sub>9</sub><sup>+</sup>], 79 (100) [C<sub>6</sub>H<sub>7</sub><sup>+</sup>], 69 (34) [C<sub>5</sub>H<sub>9</sub><sup>+</sup>].  
 30

Odor description: Musky, green, floral.


35 Example 7: Floral-musky, powdery perfume formulation for shower gel

	compound/ingredient	parts by weight 1/1000
	1. 6-Acetyl-1,1,2,4,4,7-hexamethyltetralin (Fixolide™)	68.00
	2. Ambrettolide	6.00
5	3. Benzaldehyde	0.14
	4. Citronellol, extra quality	10.00
	5. Citronellyl acetate	1.20
	6. Dipropylene glycol	135.40
	7. Elemi oil	1.20
10	8. 6-Ethyl-3-methyloct-6-en-1-ol (Super Muguet™)	2.00
	9. 4,6,6,7,8,8-Hexamethyl-1,3,4,6,7,8-hexahydrocyclopenta[g]benzopyran (Galaxolide™) 50 BB	560.00
	10. Hexyl salicylate	8.00
	11. 3-trans-Isocamphylcyclohexanol (Sandela™)	20.0
15	12. Lemon oil abergapt	10.00
	13. Linalool, synthetic	20.00
	14. Linalyl acetate, synthetic	10.00
	15. 12-Methyl-14-tetradec-9-enolide (Nirvanolide™)	2.00
	16. 15-Pentadecanolide (Thibetolide™)	4.00
20	17. 15-Pentadec-11-enolide (Habanolide™)	40.00
	18. 1-Phenylethyl acetate (Gardenol™)	1.00
	19. Rose oxide CO	0.06
	20. 1-(2,2,6-Trimethylcyclohexyl)hexan-3-ol (Timberol™)	0.20
	21. Vanillin	0.80
25	22. Propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester	100.00

The propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl  
 ester forms a very powerful and pleasant musk accord together with the polycyclic and  
 macrocyclic musk odorants, to which it adds freshness, fruitiness and a powdery  
 aspect. This accord conveys smoothness, and richness to the fragrance and imparts a  
 caressing, comfortable feeling to the perfumed product. In combination with the elemi  
 and lemon oils it as well enhances fresh and clean aspects of the perfume and makes  
 it ideally suited for application in shower gels.

Claims

1. A compound of formula (I)



(I)

wherein

R is C<sub>1</sub> to C<sub>4</sub> alkyl; or

R is vinyl or a linear, branched or cyclic C<sub>3</sub> to C<sub>4</sub> alkenyl;

X is carbonyl or a divalent radical  $-(CMe_2)-$ ;

Y is oxygen or a divalent radical  $-(CH_2)-$ ;

the bond between C-b and C-c is a single bond and the bond between C-a and C-b together with the dotted line represents a double bond; or

the bond between C-b and C-c together with the dotted line represents a double bond and the bond between C-a and C-b is a single bond.

2. A compound according to claim 1 wherein the bond between C-b and C-c together with the dotted line represents a double bond and the bond between C-a and C-b is a single bond.
3. A compound according to claim 1 wherein the bond between C-b and C-c is a single bond and the bond between C-a and C-b together with the dotted line represents a double bond.
4. A compound according to claim 1 selected from the group consisting of propanoic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, cyclopropanecarboxylic acid 2'-[1''-(3''',3'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, propionic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, cyclopropanecarboxylic acid 2'-[1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxy]-2'-methylpropyl ester, propionic acid 1''-(5''',5'''-dimethylcyclohex-1'''-

enyl)ethoxycarbonylmethyl ester, and cyclopropanecarboxylic acid 1''-(5''',5'''-dimethylcyclohex-1'''-enyl)ethoxycarbonylmethyl ester.

5. The use of a compound according to claim 1 to 4 as a fragrance.
6. The use of a compound according to claim 1 to 4 in a fragrance composition.
7. A fragrance application comprising a compound as defined in any of the claims 1 to 4, or a mixture thereof.
8. A fragrance application according to claim 7 wherein the fragrance application is a perfume, household product, laundry product, body care product or cosmetic product.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/ 3/00772

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C69/28 C07C69/74 C07C69/66 A61K7/46 C11B9/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C A61K C11B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 00/14051 A (FIRMENICH SA) 16 March 2000 (2000-03-16) cited in the application the whole document	1-8
Y	EP 0 472 966 A (FIRMENICH SA) 4 March 1992 (1992-03-04) cited in the application the whole document	1-8
P, Y	EP 1 262 474 A (GIVAUDAN SA) 4 December 2002 (2002-12-04) the whole document	1-8

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

2 March 2004

Date of mailing of the international search report

09/03/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Beslier, L



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/03/00772

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0014051	A	16-03-2000	AT 231834 T	15-02-2003
			CA 2309449 A1	16-03-2000
			CN 1275121 T	29-11-2000
			DE 69905136 D1	06-03-2003
			DE 69905136 T2	09-10-2003
			EP 1047660 A1	02-11-2000
			ES 2192067 T3	16-09-2003
			WO 0014051 A1	16-03-2000
			JP 2002524438 T	06-08-2002
<hr/>				
EP 0472966	A	04-03-1992	DE 69104311 D1	03-11-1994
			DE 69104311 T2	16-02-1995
			EP 0472966 A1	04-03-1992
			JP 2974834 B2	10-11-1999
			JP 6072952 A	15-03-1994
			US 5166412 A	24-11-1992
<hr/>				
EP 1262474	A	04-12-2002	EP 1262474 A1	04-12-2002
			WO 02096852 A1	05-12-2002
<hr/>				